

PUBLIC HEALTH FACT SHEET

Babesiosis

Massachusetts Department of Public Health, 305 South Street, Jamaica Plain, MA 02130

What is babesiosis?

Babesiosis is a disease caused by a microscopic parasite (a type of germ similar to those that cause malaria) that infects red blood cells.

Where do cases of babesiosis occur?

Babesiosis occurs in coastal areas in the northeastern United States. In Massachusetts, the disease occurs most frequently on Cape Cod and the islands of Martha's Vineyard and Nantucket, but it can occur anywhere in the state.

How is babesiosis spread?

Babesiosis is spread by the bite of an infected deer tick. The longer a tick remains attached and feeding, the higher the likelihood that it may spread the parasite. The tick must generally be attached to a person for at least 24-36 hours before it can spread the germ. Deer ticks in Massachusetts can also carry the germs that cause Lyme disease and human granulocytic anaplasmosis (also known as human granulocytic ehrlichiosis). Deer ticks are capable of spreading more than one type of germ in a single bite.

When can I get babesiosis?

Babesiosis can occur during any time of the year. The parasite that causes babesiosis is spread by infected deer ticks. Young ticks (nymphs) are most active during the warm weather months between May and July. Adult ticks are most active during the fall and spring but may also be out searching for a host any time that winter temperatures are above freezing.

How soon do symptoms of babesiosis appear after a tick bite?

Symptoms of babesiosis usually begin to appear from 1 to 8 weeks after being bitten by an infected tick.

What are the symptoms of babesiosis?

Most people who are infected by the parasite will show very mild signs of illness or no signs at all. If symptoms occur, they may include fever, chills, headache, achy joints and muscles, fatigue, nausea, vomiting, abdominal pain and dark urine. Symptoms can last for up to several months. The elderly, people without a healthy spleen, and people with weakened immune systems are more likely to develop potentially life-threatening symptoms.

Is there treatment for babesiosis?

Babesiosis can be treated with medication; however, serious complications requiring a blood transfusion and/or kidney dialysis can occur if the disease is not recognized and treated early.

What can I do to lower my chances of getting babesiosis, or any other disease, from ticks?

Prevention begins with you! Take steps to reduce your chances of being bitten by any tick. Ticks are most active during warm weather, generally late spring through fall. However, ticks can be out any time that temperatures are above freezing. Ticks cling to vegetation and are most numerous in brushy, wooded or grassy habitats. They are not found on open, sandy beaches, but may be found in grassy

dune areas. When you are outside in an area likely to have ticks (e.g. brushy, wooded or grassy places), follow these simple steps to protect yourself and your loved ones:

- Use a repellent with **DEET** (the chemical N-N-diethyl-meta-toluamide) or **permethrin** according to the instructions given on the product label. DEET products should not be used on infants under two months of age and should be used in concentrations of 30% or less on older children. Permethrin products are intended for use on items such as clothing, shoes, bed nets and camping gear, and should not be applied to skin. More information on choosing a repellent and how to use repellents safely is included in the MDPH Public Health Fact Sheet on Tick Repellents at <http://www.mass.gov/eohhs/docs/dph/cdc/factsheets/tick-repellents.pdf>. If you can't go online, contact the MDPH at (617) 983-6800 for a hard copy.
- Wear long, light-colored pants tucked into your socks or boots, and a long-sleeved shirt. This may be difficult to do when the weather is hot, but it will help keep ticks away from your skin and help you spot a tick on your clothing faster.
- Stay on cleared trails when walking or hiking, avoiding the edge habitat where ticks are likely to be.
- Talk to your veterinarian about tick control options (tick collars, repellents) for your pets.

Did you know?

You don't have to be a hiker on Cape Cod to worry about ticks. In Massachusetts, you may be bitten in your own backyard. There are lots of things you can do around your own backyard to make it less inviting for ticks! Visit the MDPH Tickborne Disease Website at <http://www.mass.gov/dph/tick> for suggestions.



After spending time in an area likely to have ticks, check yourself, your children and pets for ticks. Young ticks, called nymphs, are the size of a poppy seed. Adult deer ticks are the size of a sesame seed. Both nymph and adult deer ticks can spread the parasite that causes babesiosis; however, nymphs are more of a concern. They are aggressive feeders and so tiny that it can be difficult to see them on the body, unless you are looking carefully. When doing a tick check, remember that ticks like places that are warm and moist. Always check the back of the knees, armpits, groin, scalp, back of the neck and behind the ears. If you find a tick attached to your body, remove it as soon as possible using a fine-point tweezers. Do not squeeze or twist the tick's body, but grasp it close to your skin and pull straight out with steady pressure.

Know the symptoms of babesiosis as described in this fact sheet. If you have been someplace likely to have ticks and you develop symptoms of babesiosis, or any other disease carried by ticks, see your health care provider right away.

Where can I get more information?

- **For questions about your own health**, contact your doctor, nurse, or health care clinic.
- **For questions about babesiosis or other diseases spread by ticks**, contact the MDPH at (617) 983-6800 or toll free at (888) 658-2850 or online at <http://www.mass.gov/dph/tick>. You may also contact your local Board of Health (listed in the telephone directory under "Government").
- **For questions about tick repellents**, read the MDPH Public Health Fact Sheet on Tick Repellents at <http://www.mass.gov/eohhs/docs/dph/cdc/factsheets/tick-repellents.pdf>. If you can't go online, you may ask for hard copies of MDPH fact sheets by calling the MDPH number above.



Massachusetts Department of Public Health

Tickborne Disease Website

<http://www.mass.gov/dph/tick>



June 2012